



Palliative Care Consult Team Referral

Date of Referral (yyyy/mm/dd): Time (hh/mm):		Referral to (<i>check one</i>): ☐ Cancer Centre of South Eastern Ontario (CCSEO) – Palliative Care Clinic ☐ Belleville Cancer Center – Palliative Outreach Clinic ☐ Regional Treatment Centre Consults (Correctional Services Canada) ☐ Community Palliative Outreach Team* *Please also make a referral for homecare nursing services, if appropriate for the patient ☐ Admission assessment for Providence Care Hospital or AB Smith Homestead House (Hospice)		
Patient/Substitute Decision Maker (SDM) consents to palliative care referral (required): Yes ☐				
Last Name:		First Name:	Date of Birth (yyyy/mm/dd):	
Street:	Apartment	City/Province:	Postal Code:	
Phone:		Male ☐ Female ☐	Preferred Language:	
Substitute Decision Maker (SDM):		Contact Number:	Relationship:	
Primary Care Provider (PCP):		Phone:	Fax:	
REFERRING PROVIDER INFORMATION (ATTENDING PHYSICIAN OR NURSE PRACTITIONER)				
Printed name	Designation	Signature	Billing Number	Contact Number
Referral Type (check one): ☐ Routine (greater than 3 days) FAX 613-548-2361 ☐ Urgent (Within 3 days) Must CALL 613-548-2485 to discuss with consultant, then FAX to 613-548-2361				
Primary Diagnosis:				
Reason for Referral: (Please describe specific symptoms and medications tried to date):				
Palliative Performance Score (Required): See reverse for scoring table 10 ☐ 20 ☐ 30 ☐ 40 ☐ 50 ☐ 60 ☐ 70 ☐ 80 ☐ 90 ☐ 100 ☐				

Palliative Care Consult Team Referral

Palliative Performance Scale (PPS) v2

PPS Level	Ambulation	Activity Level & Evidence of Disease	Self -care	Intake	Conscious Level
PPS 100%	Full	Normal activity and work No evidence of disease	Full	Normal	Full
PPS 90%	Full	Normal activity and work Some evidence of disease	Full	Normal	Full
PPS 80%	Full	Normal activity and work <i>with</i> effort Some evidence of disease	Full	Normal or reduced	Full
PPS 70%	Reduced	Unable normal activity and work Significant disease	Full	Normal or reduced	Full
PPS 60%	Reduced	Unable hobby/house work Significant disease	Occasional assistance	Normal or reduced	Full or confusion
PPS 50%	Mainly sit/lie	Unable to do any work Extensive disease	Considerable assistance	Normal or reduced	Full or drowsy or confusion
PPS 40%	Mainly in bed	Unable to do most activity Extensive disease	Mainly assistance	Normal or reduced	Full or drowsy +/- confusion
PPS 30%	Totally bed bound	Unable to do any activity Extensive disease	Total care	Reduced	Full or drowsy +/- confusion
PPS 20%	Totally bed bound	Unable to do any activity Extensive disease	Total care	Minimal sips	Full or drowsy +/- confusion
PPS 10%	Totally bed bound	Unable to do any activity Extensive disease	Total care	Mouth care only	Drowsy or coma
PPS 0%	Dead				
Instructions: PPS level is determined by reading left to right to find a 'best horizontal fit.' Begin at left column reading downwards until current ambulation is determined. Then, read across to next column and downwards until each column is determined. Thus, 'leftward' columns take precedence over 'rightward' columns.					